

A new Luminous Variable in M33

A.F. Valeev,^{*} O. Sholukhova, and S. Fabrika

Special Astrophysical Observatory, Nizhnij Arkhyz 369167, Russia

30 March 2009

ABSTRACT

We present a new luminous star in M 33 located in the nuclear region. The star shows strong Fe II and [Fe II] forest, hydrogen emissions in the spectrum, as well as nebular lines. Ti II and Si II lines were detected in absorption, their radial velocity shifted by ≈ -30 km/s relative to emission lines. The star is variable over seven years with 0^m.5 variations over a year. We studied its spectral energy distribution together with five confirmed Luminous Blue Variables and Var A in M 33 using homogeneous data and methods. We found the star's bolometric luminosity to be $\log L/L_{\odot} \approx 6.27$, a surface temperature of $T \sim 16000$ K and black body temperatures of two dust components of $T \sim 900$ and 420 K. The new star has properties intermediate between the LBVs and Var A (probable cool hypergiant). In the same time it has a hot photosphere, LBV-like luminosity and an extensive circumstellar material (strong [CaII] lines). In these seven luminous variables in M 33 we find the total range of the hot component luminosities is 1.0 dex, but that of the dust componets is 2.0 dex. We conclude that the dust phenomenon in the luminous variables is temporary and variable, and that dust activity may follow strong eruptions.

Key words: Massive stars - luminous blue variables: general - stars: individual: new Luminous Variable - optical spectroscopy: stellar population - galaxies: individual - M 33

1 INTRODUCTION

Luminous Blue Variables (LBV) are the most massive stars in one of the final stages of evolution (Humphreys & Davidson 1994). They are very important for understanding of the late stages of evolution, production of WR stars, Supernovae, neutron stars and black holes. Their role in evolution of galaxies is significant. Stellar evolution calculations (e.g., Meynet et al. 2007) propose clear schemes of massive star evolution, however, the sequence of stellar conversions is not well understood yet (Smith & Owocki 2006; Smith & Conti 2008; Smith 2008). In order to understand the final evolution stages of massive stars, we need to discover and study more objects at the LBV and LBV-related stages.

Humphreys & Davidson (1994) reviewed LBV stars and their data known up to that time. In the galaxy M 33 they selected four obvious LBV objects – Var B, Var C, Var 2, Var 83, and one LBV candidate V 532 (GR 290, “Romano’s star”, (Romano 1978)). Spectroscopy has confirmed (Fabrika 2000) the LBV status of V 532. In photometric studies (Kurtev et al. 2001) its typical LBV behaviour has been confirmed as well. In following papers (Fabrika et al. 2005; Viotti et al. 2006, 2007) the star has been studied in more detail. The star Var A in this galaxy had all the properties typical for LBV stars (Hubble & Sandage 1953), it was one of the visually brightest stars in M 33 with an intermediate F-type spectrum. However, the star shown M-type spectrum (unusual for classical LBVs) after its huge brightening in 1950–1953, thus

it was classified (Humphreys et al. 1987; Humphreys et al. 2006; Viotti et al. 2006) as a cool hypergiant.

Searches for LBV candidates in M 33 have been done using various methods (Neese et al. 1991; Spiller 1992; Calzetti et al. 1995; Fabrika & Sholukhova 1995; Massey et al. 1996; Sholukhova et al. 1997; Fabrika & Sholukhova 1995; Sholukhova & Fabrika 2000; Corral & Herrero 2003). The main technique is to search for H α -emitting sources coinciding with the galaxy stars. Selections of early-type stars were preferable, however, known LBV objects show various observed colors (Sterken et al. 2008). Some works were originally addressed to a search for SS 433-like objects in M 33 (Neese et al. 1991; Calzetti et al. 1995; Fabrika & Sholukhova 1995). SS 433 has a spectrum very similar to that of a late WN star (Fabrika 2004).

A new principal step has been done by Massey et al. (2006) in their CCD survey (UBVRI, H α , [S II], [O III]) of Local Group galaxies. The final catalog in M 33 contains 146,622 stars up to 23rd magnitude with photometrical accuracy of 1–2 %. Massey et al. (2007) describe a list of H α emission-line stars in Local Group galaxies. They focused on a search for new LBVs based on their spectroscopic similarity to known LBVs. Their number of known and suspected LBVs in M 33 is 37. Some of their candidates coincide with previously described LBV candidates in M 33 (Corral 1996; Sholukhova et al. 1997, 1999; Sholukhova & Fabrika 2000; Fabrika 2000).

We performed photometry of point-like H α emitters in the M 33 images of Massey et al. (2006) and selected stars with $V < 18.5$ mag, $B - V < 0.4$ having H α excess. We did independent

^{*} E-mail: azamat@sao.ru, olga@sao.ru, fabrika@sao.ru




Figure 1. Nuclear region of M33, the bright blue object is the nucleus. The new star (J2000, 013352.42+303909 or N 93351 by Massey et al. (2006)) is indicated. The spectral slit covering both the star and another blue candidate is shown. The image was obtained as rgb from $H\alpha$, V and [O III].

aperture photometry in the UBVRI and $H\alpha$ images for selected stars. Compared to automatic PSF photometry (which is the only possible way of dealing with a huge number of stars (Massey et al. 2006)), aperture photometry is controlled by eye to avoid confusion in crowded (broad-band) and complex ($H\alpha$) regions. In the selection we used the same method as described by Fabrika et al. (1997). With a distance modulus of 24.9 mag (Bonanos et al. 2006) our list (Valeev et al. 2009) covers bright blue supergiants in the galaxy. In our follow up spectroscopy of the list stars we found a new luminous variable star. Here we report a spectral and photometrical study of the star and examine its spectral energy distribution together with all the confirmed LBVs and Var A in M33.

2 OBSERVATIONS

For spectroscopy we used the SCORPIO focal reducer (Afanasiev & Moiseev 2005) of the Russian 6-m telescope. We took spectra of the new star on November 18, 2007 with a spectral range and resolution of $\lambda\lambda 3700 - 7200 \text{ \AA}$ and $12 \text{ \AA}$, on January 7, 2008 ($\lambda\lambda 3900 - 5700 \text{ \AA}$ and $5 \text{ \AA}$), and on January 8, 2008 ($\lambda\lambda 5700 - 7400 \text{ \AA}$ and $5 \text{ \AA}$). The slit width was $1''$, and seeing $\approx 1.5''$. The spectra were reduced using standard procedures. The accuracy of radial velocity calibration is about 10 km/s .

In Fig. 1 we show the central region of M33. The star is located $34.3''$ (160 pc) from the nucleus to the South-East. The faint $H\alpha$ filaments around the star may be related to it. Our slit position was selected to cover the second blue star candidate (Fig. 1). In the HST archival broad band images the star is single and point-like.

For interstellar absorption estimates, both in this star and the other six stars in M33 (Var A, Var B, Var C, Var 2, Var 83 and V 532), we used $H\alpha/H\beta$ emission line ratios (Section 4) measured in surrounding nebulae. We used our own spectra of these targets taken with the same telescope and the MPFS instrument (IFU mode (Afanasiev et al. 2001)) in December, 2004 and October, 2005 and with SCORPIO (December, 2004 and October, 2008). The spectral data were used for independent estimates of the star temperatures.

In studying the SEDs of the stars, we used published optical photometry (Section 4) and IR archival data from 2MASS and Spitzer. The 2MASS survey of M33 is available in J, H, K bands. Spitzer observations of M33 were made using all four bands of the Infrared Array Camera (3.6, 4.5, 5.8, and $8.0 \text{ \mu m}$). We used "post-




Figure 2. Light curve of N93351 in V-band. The data for 1997–1998 were taken from Macri et al. (2001), the data for 2000–2004 from Shporer & Mazeh (2006). A systematic shift -0.2 mag was applied to the second data. Photometrical errors are less than 0.03 mag .

basic calibrated data" and did photometry of all our targets using the task *phot* in IRAF (<http://ssc.spitzer.caltech.edu/irac/dh/>). All five Spitzer observations were averaged to have one photometrical measurement for each target.

3 THE NEW LUMINOUS VARIABLE N 93351

The new star N93351 has a V magnitude of 16.16, and $B-V=0.28$, $U-B=-0.44$ (Massey et al. 2006). A light curve of the star is presented in Fig 2. The star shows a variability for the past 7 years. It shows $0^m 5$ variability over a year.

In Fig 3 we present spectrum of N93351 taken with a resolution of $5 \text{ \AA}$. The star has a blue continuum and very bright hydrogen emissions with broad wings. The most remarkable feature is Fe II and [Fe II] emission forest. We detect He I, Ti II and Si II absorption lines. He I absorption may be partly filled with nebular emission, which is seen in the spectral image. We detect Na I D1, 2 and Ca II H, K lines. The last lines may be not only interstellar in origin. The spectrum presents strong [O II] $\lambda\lambda 3727$ line, Balmer absorption edge is practically absent (Fig 4). [O III] $\lambda\lambda 4959, 5007$ lines are present, they are faint and crowded with bright Fe II emissions. There are collisionally excited nebular lines of [S II] $\lambda\lambda 6717, 6731$, [N II] $\lambda\lambda 6548, 6583$ (and marginally [N II] $\lambda 5755$) in the spectrum.

The spectrum is quite typical for LBVs. However we observe very strong [Ca II] $\lambda\lambda 7291, 7323$ emissions, which are not present in classical LBVs. These lines are seen in the cool hypergiants Var A in M33 and IRC+10420 (Humphreys et al. 1987; Humphreys et al. 2006) with extensive circumstellar material. We conclude therefore, that N93351 has intermediate spectrum between classical hot LBV stars and the cool hypergiants like Var A.

The radial velocity of Fe II and nebular lines is $\approx -120 \text{ km/s}$, which is in accordance with the H I velocity (Reakes & Newton 1978) in this area of the galaxy. We observe a blue shift of $\sim -30 \text{ km/s}$ in all absorption lines relative to hydrogen emission lines. This shift may arise in an expanding photosphere of the star. Fe II and [Fe II] line widths correspond to our spectral resolution ($5 \text{ \AA}$, measured in sky lines). For the $H\alpha$ and $H\beta$ line profiles we performed gaussian analysis for broad and narrow components and found the component widths (corrected for the instrumental resolution) of 800 km/s and 180 km/s in $H\alpha$, 950 km/s and 110 km/s in $H\beta$ correspondingly. Radial velocities of the narrow components in these two lines are about the same, $-106 \pm 5 \text{ km/s}$, and velocities of the broad components are $-125 \pm 10 \text{ km/s}$.

Table 1 presents intensities and equivalent widths of main lines, which are not blended. The intensities are given in units of intensity of $H\beta$. We present also the line intensities corrected for interstellar extinction, where we adopted $A_V = 1.0$ (Sec-




Figure 3. Spectra of N93351 in blue (top) and red (bottom) regions. Some lines are indicated (upper row), the middle row of vertical bars marks FeII lines, the bottom row marks [Fe II] lines. CCD fringes are notable at wavelengths longer than ~ 6800 Å.




Figure 4. Dereddened spectra of N93351 (middle) together with standard stars O 8I (HD 225160, top) and B 9Ib (LSI V P24, bottom) taken from Jacoby et al. (1984).

tion 4). To estimate the extinction we measured $H\alpha$ and $H\beta$ line fluxes in the surrounding nebula at a distance from $5''$ to $13''$ off the star along the slit. We found the fluxes per squared arcsecond $F(H\alpha) = (1.08 \pm 0.04) \cdot 10^{-14}$ erg/cm² sec and $F(H\beta) = (2.5 \pm 0.1) \cdot 10^{-15}$ erg/cm² sec. Corresponding line ratio $H\alpha/H\beta = 4.0$ – 4.5 gives $A_V = 1.0$ – 1.4 . The equivalent width of Na I D 1, 2 absorption (Table 1) confirms this value of extinction, a rough linear relation (Barbon et al. 1990) between the doublet EW and reddening gives $A_V = 1.2 \pm 0.2$.

Fig. 4 presents spectrum of N93351 taken with resolution 12 Å corrected for the extinction ($A_V = 1.0$, $E(B-V)=0.32$) together with standard stars O 8I ($E(B-V)=0.56$) and B 9Ib ($E(B-V)=0.23$), whose dereddened spectra were taken from Jacoby et al. (1984). Both the spectral types and reddening values of the standard stars are considered as approximate (Jacoby et al. 1984). Comparing the spectrum slope and intensity of the Balmer edge with those in classified stars we estimate roughly the temperature of N93351's photosphere of 13000 – 16000 K. Absorption Ca II K and Ti II lines may be formed in cooler extended atmosphere (the wind).

Table 1. Equivalent widths, relative intensities and dereddened ($A_V = 1.0$) relative intensities in units of $H\beta$ intensity of main not blended lines.

Line	EW ($\pm 1\sigma$), Å	I(obs)	I(dered)
[OII] $\lambda 3727$	4.9 (0.3)	0.63	0.86
CaII K (abs)	2.8 (0.3)		
H δ	1.0 (0.1)	0.11	0.13
[FeII] $\lambda 4017$	0.30 (0.02)	0.03	0.04
[FeII] $\lambda 4066$	0.33 (0.02)	0.04	0.05
H γ	2.1 (0.2)	0.18	0.21
[FeII] $\lambda 4413$	0.50 (0.02)	0.04	0.05
FeII $\lambda 4666$	0.80 (0.03)	0.07	0.07
H β	14.1 (0.1)	1.0	1.0
[FeII] $\lambda 5158$	1.1 (0.02)	0.07	0.07
FeII $\lambda 5197$	0.86 (0.03)	0.05	0.05
FeII $\lambda 5362$	0.73 (0.02)	0.04	0.04
NaI D1,D2 (abs)	1.5 (0.3)		
[FeII] $\lambda 5991$	1.70 (0.1)	0.08	0.06
FeII $\lambda 6149$	1.12 (0.05)	0.05	0.04
FeII $\lambda 6432$	1.86 (0.05)	0.09	0.06
FeII $\lambda 6516$	1.78 (0.05)	0.08	0.06
H α	124 (0.5)	5.60	3.98
[SII] $\lambda 6717 + 6731$	9.0 (0.4)	0.23	0.16
[FeII] $\lambda 7155$	3.6 (0.1)	0.12	0.08
[CaII] $\lambda 7291$	9.0 (0.1)	0.27	0.18
[CaII] $\lambda 7323$	8.9 (0.1)	0.26	0.17

4 SPECTRAL ENERGY DISTRIBUTIONS

To estimate the temperature and luminosity of this star and other six M33 stars (Var A, Var B, Var C, Var2, Var83 and V532) we study their SEDs. Our main sources for the data are optical photometry by Massey et al. (2006) (October 2000 – September 2001), 2MASS (December, 1997), Spitzer (January 2004 – June 2005) and our spectroscopy (December 2004 and October 2005). We use these data, otherwise describe individual cases. We checked all these stars in the literature for their variability so as not to use data related to different photometrical states.

We used our own spectra in order to have independent estimates of a reddening and a temperature of the stars. In the estimates

Table 2. Parameters of the studied stars. Columns present the star name, adopted values of extinction, derived bolometric luminosity, temperature of the stars (T_s), and temperatures of warm (T_w) and cold dust (T_c) components. All temperatures are given in units of 1000 K, luminosities in solar units. Contributions of the dust components (per cent) to the total luminosities are given in brackets. The star (L_s) and dust component (L_d) luminosities are given in last columns.

Name	A_V	lg L	T_s	T_w	T_c	lg L_s	lg L_d
N 93351	0.75	6.13	13	0.7(3.4)	0.19(26)	5.98	5.60
	1.0	6.27	16	0.9(0.9)	0.42(3.2)	6.25	4.88
Var A	2.0	5.63	8	1.0(15)	0.22(50)	5.17	5.44
	2.5	5.98	10	0.90(7)	0.18(64)	5.45	5.83
Var B	1.0	6.21	25	1.5(0.3)	0.18(2.5)	6.20	4.66
Var C	0.4	5.86	15	1.2(1.1)	-	5.86	3.90
Var 2	0.8	6.34	40	-	-	6.34	-
Var 83	0.7	6.32	18	1.5(0.3)	-	6.32	3.80
	1.0	6.65	22	1.5(0.1)	-	6.65	3.65
V532	0.8	6.24	35	1.5(0.5)	-	6.24	3.94

of the reddening we used $H\alpha/H\beta$ flux ratios measured in nebulae surrounding the stars. In the Case B of a standard gaseous nebula this ratio is 2.87 ($H\gamma/H\beta = 0.47$, $H\delta/H\beta = 0.26$), and stays the same to within 10% in a wide range of gas temperatures and densities (Osterbrock & Ferland 2006). Obvious limitations of this method are that the nebula may not be connected with the star, and the star may have some additional circumstellar extinction. In two stars Var A and Var C there is no clear morphological association with the nebula.

We used a standard method to fit observed SED components to black body (BB) spectra with a reddening (we adopt $R_V = 3.1$). Using the best fit SED solutions we tried to satisfy the independently estimated stellar temperatures (first) and the extinction values (second). We fitted the observed SEDs of the stars to BB spectra, suggesting that one hot spectrum is stellar and one or two cooler spectra for dust emission. We kept the temperature of cool components to less than 1500 K. We estimated luminosities integrating the extinction corrected SED components.

In some cases we did not use U data, when this band fluxes were obviously weak (because of the Balmer absorption). As a rule we did not use R, I (and sometimes J) data, because they can contain a contribution of free-free emission from stellar winds. Fig. 5 presents the observed and intrinsic SEDs of the stars as well as the best fits of BB models. Derived parameters of the stars are presented in Table 2. Below we describe the individual stars in more detail.

N 93351. The star does not show strong variability. Regarding the probable temperature and extinction ranges of the star found in Section (3), we find the best fit with $A_V \approx 1.0$ and $T \approx 16000$ K (Fig. 5). In Table 2 we present a second fit version with $A_V = 0.75$ and $T = 12800$ K, however we consider the higher temperature as more probable. The IR SED can not be fitted with one BB. We derive two dust components, with $T \approx 900$ K and $T \approx 420$ K.

Var A. In the period 1997 – 2004 the star did not change its brightness notably (Humphreys et al. 2006). From our spectrum we estimate a possible temperature range as $7000 < T < 15000$ K. Actually there are only H I emissions and absorption blends of Fe II in the spectrum. A nearby nebula located in $9''$ – $12''$ to NE direction, gives $A_V = 0.9 - 1.1$. This is the only star in our list whose luminosity was determined quite firmly (Humphreys et al.




Figure 5. Spectral energy distributions of the studied stars in M33. There are observed data (squares), data corrected for interstellar absorption (crosses), the best fit BB models of hot and dust components (dashed lines) and sum of the spectral components (solid lines).

1987; Humphreys et al. 2006), $\lg L/L_\odot \approx 6.0$ (with the distance to M33 adopted by us). These authors argue that the star has a notable amount of neutral extinction. To be consistent we use the same method as we do for other stars, but we consider this analysis as formal. We find that the luminosity $\lg L/L_\odot = 6.0$ may be reached at $A_V = 2.5$ ($T=10000$ K). One needs better spectrum of Var A to find correct extinction and the star temperature.

Var B. The star shows strong variability. After its maximum of $V \sim 15$ mag in 1992 it weakened to 17.7 mag in 2005 (Szeifert et al. 1996; Massey et al. 2006; Viotti et al. 2006; Shporer & Mazeh 2006). We used only UBV data from Viotti et al. (2006) and Spitzer data as about simultaneous. A surrounding nebula gives us the extinction limits $A_V = 1.4 - 2.8$. We estimate the star's temperature range as 15000 – 25000 K (strong He I, [Fe II], [Fe III] lines, and no Bowen blend or He II). Fitting the SED, we found the maximal value of A_V we may adopt to be 1.0 mag (corresponding to a temperature of 25000 K). Otherwise the temperature is too high. Note that in another brightness state of the star Szeifert et al. (1996) used $A_V = 0.7$. They estimated extinction as foreground galactic plus 1/2 of the value estimated from H I measurements. In spite of the different methods the star luminosity they determined is the same as that we found.

Var C. In 1985 the star showed a maximum brightness of $B=15.4$, and in 1997–2001 a minimum of $V \approx 17.3$ with a variability of less than 0.2 mag, although it begun to brighten again until 2004 on ≈ 1 mag (Szeifert et al. 1996; Shporer & Mazeh 2006). We used our main sources of the data. A nebula measured to SE direction from the star to $11''$ gives in different parts A_V from 0.32 to 0.73. The star temperature estimate is 10000 – 15000 K. There are Mg II lines, bright hydrogen emissions, and no He I lines. Fitting of the SED gives best values of $T = 14000 - 16000$ K and $A_V = 0.4$. At $A_V > 0.45$ the temperature becomes higher than we may accept from our spectrum. Note Szeifert et al. (1996) used $A_V = 0.8$, and their luminosity derived is 10% higher.




Figure 6. Temperature – luminosity diagram for seven studied M33 stars. In N93351 and Var83 we used two most probable values of extinction, inner area shows the acceptable L, T values. In VarA our analysis is rather formal (see text). In other stars we used one value of interstellar extinction (Table 2). The segments indicate a range of acceptable temperatures. Stellar evolution tracks (Claret 2004) for $Z=0.007$ stars are plotted.

Var83. The star did not show strong variability in 1997 – 2004 (Mochejska et al. 2001; Massey et al. 2006); its variations were less than 0.5 mag. Its nebula indicates $A_V \approx 1.35$. In our spectrum there are very strong He I, Fe II, [Fe II], [Fe III] and Si II emissions (no Bowen blend), which implies a temperature range 15000 – 25000 K. Our SED solutions give $T > 30000$ K at $A_V \approx 1.35$, which is in contradiction with the spectrum. Hence we use two values $A_V = 0.7$ (Fig. 5) and 1.0 giving the most close temperatures to those estimated from the spectrum. Note Szeifert et al. (1996) used $A_V = 1.05$, they obtained the star luminosity inside the range of our limits Table 2.

Var2. The star has not shown strong variability since 1935 (< 0.3 mag), now its brightness is about $V \approx 18.2$ (Szeifert et al. 1996; Massey et al. 2006). There are no 2MASS and Spitzer data (the star is faint). We show the K band point in Fig. 5 from Szeifert et al. (1996) as an upper limit. We found $A_V \approx 0.85$ in Var2's nebula. There are He I, He II and Bowen blend lines in Var2 spectrum. The temperature is estimated as 35000 – 50000 K. The best fit of the SED gives estimates $A_V \approx 0.8$ and $T \approx 40000$ K.

V532. The star shows strong historical variability, however in the period 1997 – 2005 it was relatively calm (Kurtev et al. 2001; Zharova & Sholukhova 2004; Viotti et al. 2007, and our own unpublished data), from $B \approx 17.2$ in 1997 to 17.4 in 2000, dropping to 17.7 to October 2001, and returning to 17.4 in 2004 – beginning of 2005. A close nebula surrounding V532 indicates $A_V \sim 1.1$. There are He II, Bowen blend and strong He I emissions in our spectra, which implies that the temperature has to be 30000 – 40000 K. The best fit of the SED we find with $T = 35000$ K ($A_V = 0.8$).

5 CONCLUDING REMARKS

Fig. 6 shows the temperature – luminosity diagram for the seven studied stars in M33. Stellar evolution tracks (Claret 2004) for 40 – 125 $M_\odot$ stars are plotted, which were abridged in the latest evolution stages so as not to crowd the figure. All these stars have masses of $\sim 40 - 130 M_\odot$, and are located in the region of evolved LBV stars. The new star N93351 is among the brightest evolved stars in M33.

One finds from Table 2 that the dust reradiates on average $\sim 8\%$ of the energy budget of the stars. The total scatter of stellar component luminosities is 1.0 dex, while that of the dust luminosities is 2.0 dex. Any possible uncertainties in interstellar extinction are not as important in IR radiation as they are in the optical range, but we observe much bigger scatter in the dust component luminosities than that in stellar components. This may mean that the dust phenomenon in the most massive evolved stars is quite temporary, and that the dust components must be variable on scales much less than those of core evolution. One may conclude that the dust activity of LBV and related stars follows the strong eruptions (at time scale of tens, or possibly, hundreds of years), as was detected in VarA in 1950–1953 (Humphreys et al. 1987).

The new star N93351 has properties intermediate between the LBVs and VarA, in the same time it has a hot photosphere, LBV-like luminosity and an extensive circumstellar material. This makes N93351 very important for detail studies in future.

ACKNOWLEDGMENTS

The authors thank E. Chentsov for helpful discussions, the referee R. Humphreys for very important comments, P. Massey for providing us with M33 photometrical data before publication, O. Maryeva for help in spectral data reduction, P. Boley for correction of the manuscript. This work is based in part on archival data obtained with the Spitzer Space Telescope, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. The research was supported by RFBR grants N 06-02-16865, 07-02-00909.

REFERENCES

- Afanasiev, V. L., Dodonov, S. N., Moiseev, A. V., 2001, in "Stellar Dynamics: From Classic to Modern". L. P. Ossipkov and I. I. Nikiforov (eds.). Saint Petersburg: Sobolev Astronomical Institute, 103
- Afanasiev, V. L., Moiseev, A. V., 2005, *AstL*, 31, 194
- Barbon, R., Benetti, S., Rosino, L., Cappellaro, E., Turatto, M., 1990, *A&A*, 237, 79
- Bonanos, A. Z., Stanek, K. Z., Kudritzki, R. P., et al., 2006, *ApJ*, 652, 313
- Calzetti, D., Kinney, A. L., Ford, H., Doggett, J., Long, K. S., 1995, *AJ*, 110, 2739
- Claret, A., 2004, *A&A*, 453, 769
- Corral, L. J., 1996, *AJ*, 112, 1450
- Corral, L. J., Herrero, A., 2003, *RMxAA*, 16, 265
- Fabrika, S., 2000, in "Thermal and Ionization Aspects of Flows from Hot Stars: Observations and Theory", H.J.G.L.M Lamers and A. Sagar (eds), ASP Conf, 204, 57.
- Fabrika, S., 2004, *Astrophysics and Space Physics Reviews*, 12, 1
- Fabrika, S., Sholukhova, O. 1995, *Ap&SS*, 226, 229

- Fabrika, S. N., Sholukhova, O. N., Zakharova, S. A., 1997, BSAO, 43, 133
- Fabrika, S., Sholukhova, O. 1999, A&AS, 140, 309
- Fabrika, S., Sholukhova, O., Becker, T., Afanasiev, V., Roth, M., Sanchez, S. F., 2005, A&A, 437, 217
- Hubble, E., Sandage, A., 1953, ApJ, 118, 353
- Humphreys, R. M., Blaha, C., D’Odorico, S., Gull, T. R., Benvenuti, P., 1984, 278, 124
- Humphreys, R. M., Davidson, K., 1994, PASP, 106, 1025
- Humphreys, R. M., Jones, T. J., Gehrz, R. D., 1987, AJ, 94, 315
- Humphreys, R. M., Jones, T. J., Polomski, E., et al., 2006, AJ, 131, 2105
- Jacoby, G. H., Hunter, D. A., Christian, C.A., 1984, ApJS, 56, 257
- Kurtev, R., Sholukhova, O., Borissova, J., Georgiev, L., 2001, RMxAA, 37, 57
- Macri, L. M., Stanek, K. Z., Sasselov, D. D., Krockenberger, M., Kaluzny, J., 2001, AJ, 121, 870
- Massey, P., Bianchi, L., Hutchings, J. B., Stecher, T. P., 1996, ApJ, 469, 629
- Massey, P., Olsen, K. A. G., Hodge, P. W., Strong, S. B., Jacoby, G. H., Schlingman, W., Smith, R. C., 2006, AJ, 131, 2478
- Massey, P., McNeill, R. T., Olsen, K. A. G., Hodge, P. W., Blaha, C., Jacoby, G. H., Smith, R. C., Strong, S. B., 2007, AJ, 134, 2474
- Meynet, G., Eggenberger, P., Maeder, A., 2007, in "Populations of massive stars in galaxies, implications for the stellar evolution theory", IAU Symp. N 241, A. Vazdekis and R. F. Peletier (eds.), 13
- Mochejska, B. J., Kaluzny, J., Stanek, K. Z., Sasselov, D. D., Szentgyorgyi, A. H., 2001, AJ, 122, 2477
- Neese, C. L., Armandroff, T. E., Massey, P., 1991, in "Wolf-Rayet Stars and Interrelations with Other Massive Stars in Galaxies", IAU Symp. N 143, K. A. van der Hucht & B. Hidayat (eds.), 651
- Osterbrock D. E., Ferland G. J., 2006, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei, Sausalito, CA: University Science Books
- Reakes, M. L., Newton, K. 1978, MNRAS, 185, 277
- Romano, G., 1978, A&A, 67, 291
- Sholukhova, O. N., Fabrika, S. N., Vlasyuk, V. V., Burenkov, A. N., 1997, AstL, 23, 458
- Sholukhova, O. N., Fabrika, S. N., Vlasyuk, V. V., 1999, AstL, 25, 14
- Sholukhova, O., Fabrika, S., 2000, in "Stars, Gas and Dust in Galaxies: Exploring the Links", ASP Conf., v. 221, D. Alloin, K. Olsen, and G. Galaz (eds.), 171
- Shporer, A., Mazeh, T., 2006, MNRAS, 370, 1429
- Smith, N., Owocki, S. P., 2006, ApJ, 645, L45
- Smith, N., 2008, in "Mass Loss from Stars and the Evolution of Stellar Clusters", ASP Conf., v. 388, A. de Koter, L. J. Smith, and L. B. F. M. Waters (eds.), 129
- Smith, N., Conti, P., 2008, ApJ, 679, 1467
- Spiller, F., 1992, PhDT,
- Sterken, C., van Genderen, A. M., Plummer, A., Jones, A. F., 2008, A&A, 484, 463
- Szeifert, T., Humphreys R. M., Davidson, K., Jones, T. J., Stahl, O., Wolf, B., Zickgraf, F.-J., 1996, A&A, 314, 131
- Valeev, A. F., Sholukhova, O. N., Fabrika, S. N., 2009, in preparation.
- Viotti, R. F., Rossi, C., Polcaro, V. F., Montagni, F., Gualandi, R., Norci, L., 2006, A&A, 458, 225
- Viotti, R. F., Galleti, S., Gualandi, R., Montagni, F., Polcaro, V. F., Rossi, C., Norci, L., 2007, A&A, 464, L53
- Zharova, A., Sholukhova, O., 2004, CoAst, 145, 28